

Other relevant publications on papilionoid legumes

- Barker, N. P., Schrire, B. D., & Kim, J.-H. 2000. Generic relationships in the tribe Indigofereae (Leguminosae: Papilionoideae) based on sequence data and morphology. Pp. 311-337, in Herendeen, P. S., & Bruneau, A. (eds), *Advances in Legume Systematics, Part 9*. Royal Botanic Gardens, Kew.
- Boatwright, J. S., Savolainen, V., Van Wyk, B.-E., Schutte-Vlok, A. L., Forest, F., & Van der Bank, M., 2008a. Systematic position of the anomalous genus *Cadia* and the phylogeny of the tribe Podalyrieae (Fabaceae). *Syst. Bot.* 33: 133-147.
- Boatwright, J. S., Le Roux, M. M., Wink, M., Morozova, T., & Van Wyk, B. E., 2008. Phylogenetic relationships of tribe Crotalarieae (Fabaceae) inferred from DNA sequences and morphology. *Syst. Bot.* 33: 752-761.
- Boatwright, J. S., Wink, M., & Van Wyk, B.-E., 2011. The generic concept of *Lotononis* (Crotalarieae, Fabaceae): reinstatement of the genera *Euchlora*, *Leobordea* and *Listia* and the new genus *Ezoloba*. *Taxon* 60: 161-177.
- Castellanos, C., Steeves, R., Lewis, G. P., & Bruneau, A. 2017. A settled sub-family for the orphan tree: The phylogenetic position of the endemic Colombian genus *Orphanodendron* in the Leguminosae. *Brittonia* 69: 62-70.
- Cooper, W., Crayn, D. M., Zich, F. A., Miller, R. E., Harrison, M., & Nauheimer, L. 2019. A review of *Austrocallerya* and *Pongamia* (Leguminosae subfamily Papilionoideae) in Australia, and the description of a new monotypic genus, *Ibatiria*. *Australian Syst. Bot.* 32: 363-384.
- Crisp, M. D., Gilmore, S., & van Wyk, B.-E. 2000. Molecular phylogeny of the genistoid tribes of papilionoid legumes. Pp. 249-276, in Herendeen, P. S., & Bruneau, A. (eds), *Advances in Legume Systematics, Part 9*. Royal Botanic Gardens, Kew.
- Crisp, M. D., & Cook, L. G. 2003. Phylogeny and embryo sac evolution in the endemic Australasian papilionoid tribes Mirbelieae and Bossiaeeae. Pp. 253-268, in Klitgaard, B. B. & Bruneau, A. (eds), *Advances in Legume Systematics, Part 10, Higher Level Systematics*. Royal Botanic Gardens, Kew.
- Degtjareva, G. V., Valiejo-Roman, C. M., Samigullin, T. H., Guara-Requena, M., & Sokoloff, D. D. 2012. Phylogenetics of Anthyllis (Leguminosae: Papilionoideae: Loteae): Partial incongruence between nuclear and plastid markers; A long branch problem and implications for morphological evolution. *Mol. Phylogenet. Evol.* 62: 693-707.
- De Moura, T. M., Wilmot-Dear, M., Vatanparast, M., Fortuna-Perez, A. P., Tozzi, A. M. G. A., & Lewis, G. P. 2016. A new infrageneric classification of *Mucuna* (Leguminosae-Papilionoideae) supported by morphology, molecular phylogeny and biogeography. *Syst. Bot.* 41: 606-616.
- Diez, M. J., & Ferguson, I. K. 1996. Studies on the pollen morphology and taxonomy of the tribes Loteae and Coronilleae (Papilionoideae; Leguminosae) 3, *Coronilla* L. and related genera and systematic conclusions. *Review Palaeobot. Palynol.* 94: 239-257.
- Doyle JJ, Doyle JL, Ballenger JA, Dickson EE, Kajita T, Ohashi H. 1997. A phylogeny of the chloroplast gene *rbcL* in the Leguminosae: taxonomic correlations and insights into the evolution of nodulation. *Amer J Bot* 84: 541-554.
- Doyle JJ, Chappill JA, Bailey, CD, Kajita T. 2000. Towards a comprehensive phylogeny of legumes: Evidence from *rbcL* sequences and non-molecular data. In P. S.

- Herendeen and A. Bruneau (eds.), *Advances in legume systematics*, part 9, 1–20. Royal Botanic Gardens, Kew, UK.
- Duan, L., Yang, X., Liu, P.-L., Arsian, E., Ertuğrul, K., & Chang, Z.-Y. 2015. Phylogeny of *Hedysarum* and tribe Hedysareae (Leguminosae: Papilionoideae) inferred from sequence data of ITS, *matK*, *trnL-F* and *psbA-trnH*. *Taxon* 64: 49-64.
- Duan, L., Yang, X., Liu, P.-L., Johnson, G., Wen, J., & Chang, Z. 2016. A molecular phylogeny of *Caragana* (Leguminosae, Papilionoideae) reveals insights into new generic and infrageneric delimitations. *Phytokeys* 70: 111-137.
- Duan, L. [et al. 2019], Harris, A. J., Su, C., Yw, W., Deng, S.-W., Fu, L., Wen, J., & Chen, H.-F. 2020. A fossil-calibrated phylogeny reveals the biogeographic history of the *Cladrastis* clade, an amphi-Pacific early-branching group in papilionoid legumes. *Molec. Phyl. Evol.* 143:106673.
- Egan, A. N. & Crandall, K. A. 2008. Incorporating gaps as phylogenetic characters across eight DNA regions: Ramifications for North American Psoraleeae (Leguminosae). *Mol. Phylogenet. Evol.* 46: 532-546.
- Egan, A. N. & Doyle, J. 2010. A comparison of global, gene-specific, and relaxed clock methods in a comparative genomics framework: Dating the polyploid history of soybean (*Glycine max*). *Syst. Biol.* 59: 534-547.
- Egan, A. N., Vatanparast, M., & Cagle, W. 2016. Parsing polyphyletic *Pueraria*: delimiting distinct evolutionary lineages through phylogeny. *Mol. Phylogenet. Evol.* 104: 44-59.
- Hughes, C. E., Lewis, G. P., Daza, Y. A., & Reynel, C. 2004. *Maraniona*. A new dalbergioid legume genus (Leguminosae: Papilionoideae) from Peru. *Syst. Bot.* 29: 366-374.
- Hughes, C. E. & Eastwood, R. 2006. Island radiation on a continental scale: Exceptional rates of plant diversification after uplift of the Andes. *Proc. Natl. Acad. Sci. U.S.A.* 103: 10334-10339.
- Ireland, H. E., Kite, G. C., Veitch, N. C., Chase, M. W., Schrire, B., Lavin, M., Linares, J., & Pennington, R. T. 2010. Biogeographical, ecological and morphological structure in a phylogenetic analysis of *Ateleia* (Swartzieae, Fabaceae) derived from combined molecular, morphological and chemical data. *Bot. J. Linn. Soc.* 162: 39-53.
- Jabbour, F., Gaudeul, M., Lambourdière, J., Ramstein, G., Hassanin, A., Labat, J. N., & Sarthou, C. 2018 [= 2017]. Phylogeny, biogeography and character evolution in the tribe Desmodieae (Fabaceae: Papilionoideae), with special emphasis on the New Caledonian endemic genera. *Molec. Phyl. Evol.* 118: 108-121.
- Kajita, T., Ohashi, H., Tateishi, Y., Bailey, C. D., & Doyle, J. J., 2001. *rbcL* and legume phylogeny, with particular reference to Phaseoleae, Millettieae, and allies. *Syst. Bot.* 26: 515-536.
- Käss E, Wink M. 1995. Molecular phylogeny of the Papilionoideae (family Leguminosae): *rbcL* gene sequences versus chemical taxonomy. *Bot. Acta* 108: 149–162 .
- Käss E, Wink M. 1996. Molecular evolution of the Leguminosae: Phylogeny of the three subfamilies based on *rbcL* sequences. *Biochem Syst Ecol* 24: 365–378.

- Käss, E., & Wink, M. 1997. Phylogenetic relationships in the Papilionoideae (family Leguminosae) based on nucleotide sequences of cpDNA (*rbcL*) and ncDNA (ITS1 and 2). *Molec. Phyl. Evol.* 8: 65-88.
- Klitgård, B. B., Forest, F., Booth, T. J., & Saslis-Lagoudakis, C. H. 2013. A detailed investigation of the Pterocarpus clade (Leguminosae: Dalbergieae): *Etaballia* with radially symmetrical flowers is nested within the papilionoid-flowered *Pterocarpus*. *S. Afr. J. Bot.* 89: 128-142.
- Koenen, E. J., Ojeda, D. I., Bakker, F. T., Wieringa, J. J., Kidner, C., Hardy, O. J., ... & Hughes, C. E. (2020). The origin of the legumes is a complex paleopolyploid phylogenomic tangle closely associated with the Cretaceous-Paleogene (K-Pg) mass extinction event. *Systematic Biology*. Accepted manuscript.
- Koenen, E. J., Ojeda, D. I., Steeves, R., Migliore, J., Bakker, F. T., Wieringa, J. J., ... & Hughes, C. E. (2020). Large-scale genomic sequence data resolve the deepest divergences in the legume phylogeny and support a near-simultaneous evolutionary origin of all six subfamilies. *New Phytologist*, 225(3), 1355-1369.
- Lavin, M., & Sousa M., S. 1995. Phylogenetic systematics and biogeography of the tribe Robinieae. *Syst. Bot. Monogr.* 45: 1-165.
- Lavin, M., Doyle, J. J., & Palmer, J. D. 1990. Evolutionary significance of the loss of the chloroplast-DNA inverted repeat in the Leguminosae subfamily Papilionoideae. *Evolution* 44: 390-402.
- Lavin, M., Eshbaugh, E., Hu, J., Mathews, S., & Sharrock, R. A. 1998. Monophyletic subgroups of the tribe Millettieae (Leguminosae) as revealed by phytochrome nucleotide sequence data. *American J. Bot.* 85: 412-433.
- Lavin, M., Thulin, M., Labat, J.-N., & Pennington, R. T. 2000. Africa, the odd man out: Molecular biogeography of dalbergioid legumes (Fabaceae) suggests otherwise. *Syst. Bot.* 25: 449-467.
- Lavin, M., Pennington, R. T., Klitgaard, B. B., Sprent, J. I., de Lima, H. C., & Gasson, P. E. 2001. The dalbergioid legume (Fabaceae): Delimitation of a pantropical monophyletic clade. *American J. Bot.* 88: 503-533.
- Le Roux, M. M., Boatwright, J. S., & Van Wyk B.-E. A global infrageneric classification system for the genus *Crotalaria* (Leguminosae) based on molecular and morphological evidence. *Taxon* 62: 957-71.
- Lewis G, Schrire B, Mackinder B, Lock M. 2005. Legumes of the World. Royal Botanic Gardens, Kew, UK.
- Li, Q.-G., Li, Z., Li, C., Dunwell, J. M., & Zhang, Y.-M. 2013. Comparative genomics suggests that an ancestral polyploidy event leads to enhanced root nodule symbiosis in the Papilionoideae. *Mol. Biol. Evol.* 30: 2602-2611.
- McMahon, M. M. 2005. Phylogenetic relationships and floral evolution in the papilionoid legume clade Amorpheae. *Brittonia* 57: 397-411.
- McMahon, M. M., & Hufford, L. 2005. Evolution and development in the amorphoid clade (Amorpheae: Papilionoideae: Leguminosae): Petal loss and dedifferentiation. *Internat. J. Plant Sci.* 166: 383-396.
- McMahon, M. M., & Hufford, L. 2006. Phylogeny of Amorpheae (Fabaceae: Papilionoideae). *Amer. J. Bot.* 91: 1219-1230.
- McMahon, M. M., & Sanderson, M. J. 2006. Phylogenetic supermatrix analysis of GenBank sequences from 2228 papilionoid legumes. *Syst. Biol.* 55: 818-836.

- Mansano, V. de F., Bittrich, V., Tozzi, A. M. G. de A., & de Souza, A. P. 2004. Composition of the *Lecointea* clade (Leguminosae, Papilionoideae, Swartzieae), a re-evaluation based on combined evidence from morphology and molecular data. *Taxon* 53: 1007-1018.
- Meireles, J. E., Azevedo-Tozzi, A. M. G., & Lavin, M. 2014. A phylogenetic analysis of molecular and morphological data reveals a paraphyletic *Poecilanthe* (Leguminosae, Papilionoideae). *Syst. Bot.* 39: 1142–1149.
- Moura, T. M., Vatanparast, M., Tozzi, A. M. G. A., Forest, F., Wilmot-Dear, C. M., Simon, M. F., Mansano, V. F., Kajita, T., & Lewis, G. P. 2016 [= 2015]. A molecular phylogeny and a new infrageneric classification of *Mucuna* Adans. (Leguminosae-Papilionoideae) including insights from morphology and hypotheses about biogeography. *Internat. J. Plant Sci.* 177: 76-89.
- Pennington, R. T., Klitgaard, B., Ireland, H., & Lavin, M. 2000. New insights into floral evolution of basal Papilionoideae from molecular phylogenies. Pp. 233-248, in Herendeen, P. S., & Bruneau, A. (eds), *Advances in Legume Systematics, Part 9*. Royal Botanic Gardens, Kew.
- Pennington RT, Lavin M, Ireland H, Klitgaard B, Preston J, Hu J-M. 2001. Phylogenetic relationships of basal papilionoid legumes based upon sequences of the chloroplast trnL intron. *Syst. Bot.* 26: 537–556.
- Pennington, R. T., Lavin, M., Särkinen, T., Lewis, G. P., Klitgaard, B. B., & Hughes, C. E. 2010. Contrasting plant diversification histories within the Andean biodiversity hotspot. *Proc. Natl. Acad. Sci. U.S.A.* 107: 13783-13787.
- Pirie, M. D., Klitgaard, B. B., & Pennington, R. T. 2009. Revision and biogeography of *Centrolobium* (Leguminosae, Papilionoideae). *Syst. Bot.* 34: 345-359.
- Rockinger, A., Flores, A. S., & Renner, S. S. 2017. Clock-dated phylogeny for 48% of the 700 species of *Crotalaria* (Fabaceae–Papilionoideae) resolves sections worldwide and implies conserved flower and leaf traits throughout its pantropical range. *BMC Evolutionary Biology* 17: 61.
- Safaei Chaei Kar, S., Ghanavati, F., Naghavi, M. R., Amirabadi-zade, H., & Rabiee, R. 2014. Molecular phylogenetics of the *Onobrychis* genus (Fabaceae: Papilionoideae) using ITS and *trnL–trnF* DNA sequence data. *Australian J. Bot.* 62: 235-250.
- Saslis-Lagoudakis, C. H., Chase, M. W., Robinson, D. N., Russell, J., & Klitgård, B. B. 2008. Phylogenetics of neotropical *Platymiscium* (Leguminosae: Dalbergieae): Systematics, divergence times, and biogeography inferred from nuclear ribosomal and plastid DNA sequence data. *Amer. J. Bot.* 95: 1270-1286.
- Schrire, B. D., Lavin, M., & Lewis, G. P. 2005. Global distribution patterns of the Leguminosae: Insights from recent phylogenies. *Biol. Skr.* 55: 375-422.
- Schrire, B. D., Lavin, M., Barker, N. P., & Forest, F. 2009. Phylogeny of the tribe Indigofereae (Leguminosae-Papilionoideae): Geographically structured more in succulent-rich and temperate settings than in grass-rich environments. *Amer. J. Bot.* 96: 816-852.
- Sirichamorn, Y., Adema, F. A. C. B., Gravendeel, B., & Van Welzen, P. C. 2012. Phylogeny of palaeotropical *Derris*-like taxa (Fabaceae) based on chloroplast and nuclear DNA sequences shows reorganization of (infra)generic classifications is needed. *Am. J. Bot.* 99: 1793-1808.

- Sirichamorn, Y., Adema, F. A. C. B., Roos, M. C., & Van Welzen, P. C. 2014. Molecular and morphological phylogenetic reconstruction reveals a new generic delimitation of Asian *Derris* (Fabaceae): reinstatement of *Solori* and synonymisation of *Paraderris* with *Derris*. *Taxon* 63: 522–538.
- Stefanovic S, Pfeil BE, Palmer JD, Doyle JJ. 2009. Relationships among phaseoloid legumes based on sequences from eight chloroplast regions. *Syst Bot* 34: 115–128.
- Torke BM, Schaal BA. 2008. Molecular phylogenetics of the species-rich neotropical genus *Swartzia* (Leguminosae, Papilionoideae) and related genera of the swartzioid clade. *Amer. J. Bot.* 95: 215–228.
- Van Wyk, B.-E. 2003. The value of chemosystematics in clarifying relationships in the genistoid tribe of papilionoid legumes. *Biochem. Syst. Ecol.* 31: 875-884.
- Vatanparast, M., Klitgård, B. B., Adema, F. A. C. B., Pennington, R. T., Yahara, T., & Kajita, T. 2013. First molecular phylogeny of the pantropical genus *Dalbergia*: implications for infrageneric circumscription and biogeography. *S. Afr. J. Bot.* 89: 143–149.
- Vatanparast M, Powell A, Doyle JJ, Egan AN. 2018. Targeting legume loci: A comparison of three methods for target enrichment bait design in Leguminosae phylogenomics. *Appl. Plt. Sci.* 6(3):e103.