

Previous publications from participants on papilionoid legumes

- Cai Z, Guisinger M, Kim H-G, Ruck E, Blazier JC, McMurtry V, Kuehl JV, Boore J, Jansen RK 2008. Extensive reorganization of the plastid genome of *Trifolium subterraneum* (Fabaceae) is associated with numerous repeated sequences and novel DNA insertions. *J Mol Evol* 67: 696–704.
- Cardoso, D., Lima, H. C. de, Rodrigues, R. S., Queiroz, L. P. de, Pennington, R. T., & Lavin, M. 2012a. The realignment of *Acosmium* sensu stricto with the Dalbergioid clade (Leguminosae: Papilionoideae) reveals a proneness for independent evolution of radial floral symmetry among early-branching papilionoid legumes. *Taxon* 61: 1057-1073.
- Cardoso, D., Lima, H. C. de, Rodrigues, R. S., Queiroz, L. P. de, Pennington, R. T., & Lavin, M. 2012b. The *Bowdichia* clade of genistoid legumes: Phylogenetic analysis of combined molecular and morphological data and a recircumscription of *Diploptropis*. *Taxon* 61: 1074-1087.
- Cardoso, D, Queiroz, L. P. de, Pennington, R. T., Lima, H.C. de, Fonty, E., Wojciechowski, M.F., & Lavin, M. 2012c. Revisiting the phylogeny of papilionoid legumes: new insights from comprehensively sampled early-branching lineages. *Amer. J. Bot.* 99: 1991-2013.
- Cardoso, D., Queiroz, L. P. de, Lima, H. C. de, Sukanuma, E., van den Berg, C., & Lavin, M. 2013a. A molecular phylogeny of the vataireoid legumes underscores floral evolvability that is general to many early-branching papilionoid lineages. *Amer. J. Bot.* 100: 403-421.
- Cardoso D., Pennington, R. T., Queiroz L. P. de, Boatwright, J.S., van Wyk, B.-E., Wojciechowski, M. F., & Lavin, M. 2013b. Reconstructing the deep-branching relationships of the papilionoid legumes. *S. African J. Bot.* 89: 58-75.
- Cardoso, D., São-Mateus, W. M., Cruz, D. T., Zartman, C. E., Komura, D. L., Kite, G., Prenner, G., Wieringa, J. J., Clark, A., Lewis, G., Pennington, R. T., Queiroz, L. P., 2015. Filling in the gaps of the papilionoid legume phylogeny: the enigmatic Amazonian genus *Petaladenium* is a new branch of the early-diverging Amburaneae clade. *Mol. Phylogenet. Evol.* 84: 112-124.
- Champagne, C. E. M., Goliber, T. E., Wojciechowski, M. F., Mei, R. W., Townsley, B. T., Wang, K., Paz, M. M., Geeta, R., & Sinha, N. R. 2007. Compound leaf development and evolution in the legumes. *Pl. Cell* 19: 3369-3378.
- Cardoso, D., Harris, D. J., Wieringa, J. J., São-Mateus, W. M. B., Batalha-Filho, H., Torke, B. M., Prenner, G., & Queiroz, L. P. de. 2017. A molecular-dated phylogeny and biogeography of the monotypic legume genus *Haplormosia*, a missing African branch of the otherwise American-Australian Brongniartieae clade. *Mol. Phylogenet. Evol.* 107: 431-442.
- Choi, I. S., and Choi, B. H. 2017. The distinct plastid genome structure of *Maackia fauriei* (Fabaceae: Papilionoideae) and its systematic implications for genistoids and tribe Sophoreae. *PLoS One*, 12: e0173766.
- Choi I-S, Jansen RK, Ruhlman TA. 2019. Lost and found: Return of the inverted repeat in the legume clade defined by its absence. *Gen Biol Evol* 11: 1321–1333.

- Choi I-S, Schwarz EN, Ruhlman TA, Khiyami MA, Sabir JMS, Hajarrah NH, Sabir MJ, Rabah SO, Jansen RK. 2019. Fluctuations in Fabaceae mitochondrial genome size and content are both ancient and recent. *BMC Plt Biol* 19: 448.
- Choi I.-S., Ruhlman T.A., and Jansen R.K. 2020. Comparative mitogenome analysis of the genus *Trifolium* reveals independent gene fission of ccmFn and intracellular gene transfers in Fabaceae. *International Journal of Molecular Sciences* 21, 1959.
- Cruz, D. T., Idárraga, A., Banda, K., Cogollo, A., van den Berg, C., Queiroz, L. P. de, Pennington, R. T., Lavin, M. & Cardoso, D. 2018. Ancient speciation of the papilionoid legume *Luetzelburgia jacana*, a newly discovered species in an inter-Andean seasonally dry valley of Colombia. *Taxon* 67: 931-943.
- Da Silva, M. J., de Queiroz, L. P., Tozzi, A. M. G. de A., Lewis, G. P., & de Sousa, A. P. 2012. Phylogeny and biogeography of *Lonchocarpus* sensu lato and its allies in the tribe Millettieae (Leguminosae, Papilionoideae). *Taxon* 61: 93-108.
- De Queiroz, L. P., Pastore, J. F. B., Cardoso, D., Snak, C., Lima, A. L. de C., Gagnon, E., Vatanparast, M., Holland, A. E., & Egan, A. N. 2015. A multilocus phylogenetic analysis reveals the monophyly of a recircumscribed papilionoid legume tribe Diocleae with well-supported generic relationships. *Molec. Phyl. Evol.* 90: 1-19.
- De Queiroz, L. P., São-Mateus, W., Delgado-Salinas, A., Torke, B. M., Lewis, G. P., Dorado, O., Ardley, J. J., Wojciechowski, M. F., & Cardoso, D. 2017. A molecular phylogeny reveals the Cuban enigmatic genus *Behaimia* as a new piece in the Brongniatiaceae puzzle of papilionoid legumes. *Molec. Phyl. Evol.* 109: 191-202.
- Fortuna-Perez, A. P., Silva, M. J., Queiroz, L. P., Lewis, G. P., Simões, A. O., Tozzi, A. M. G. A., Särkinen, T. E., & Souza, A. P. 2013. Phylogeny and biogeography of the genus *Zornia* (Leguminosae: Papilionoideae: Dalbergieae). *Taxon* 62: 723-732.
- Hu, J-M., M. Lavin, M. F. Wojciechowski, and M. J. Sanderson. 2000. Phylogenetic systematics of the tribe Millettieae (Leguminosae) based on matK sequences, and implications for evolutionary patterns in Papilionoideae. *American J. Botany* 87: 418-430.
- Hu, J-M., M. Lavin, M. F. Wojciechowski, and M. J. Sanderson. 2002. Phylogenetic analysis of nuclear ribosomal ITS/5.8 S sequences in the tribe Millettieae (Fabaceae): *Poecilanthe-Cyclobium*, the core Millettieae, and the *Callerya* group. *Syst. Bot.* 27: 722-733.
- Jansen RK, Wojciechowski MF, Sanniyasi E, Lee S-B, Daniell H. 2008. Complete plastid genome sequence of the chickpea (*Cicer arietinum*) and the phylogenetic distribution of *rps12* and *clpP* intron losses among legumes (Leguminosae). *Mol Phylogen Evol* 48: 1204–1217.
- Jin, D. P., Choi, I. S., and Choi, B. H. 2019. Plastid genome evolution in tribe Desmodieae (Fabaceae: Papilionoideae). *PloS one*, 14(6): e0218743.
- Koenen, E., De Vos, J. M., Atchison, G., Simon, M. F., Schrire, B., Souza, E. R., Queiroz, L. P., & Hughes, C. E. 2013. Exploring the tempo of species diversification in legumes. *S. African J. Bot.* 89: 19-30.
- Lavin, M., Pennington, R. T., Klitgaard, B. B., Sprent, J. I., Lima, H. C., & Gasson, P. E., 2001. The dalbergioid legumes (Fabaceae): delimitation of a pantropical monophyletic clade. *Am. J. Bot.* 88: 503-533.
- Lavin, M., Wojciechowski, M. F., Gasson, P., Hughes, C. & Wheeler, E.. 2003. Phylogeny of robinoid legumes (Fabaceae) revisited: *Coursetia* and *Gliricidia*

- recircumscribed, and a biogeographical appraisal of the Caribbean endemics. *Syst. Bot.* 28: 387-409.
- Lavin, M., Schrire, B. P., Lewis, G., Pennington, R. T., Delgado-Salinas, A., Thulin, M., Hughes, C. E., Beyra-Matos, A., & Wojciechowski, M. F. 2004. Metacommunity process rather than continental tectonic history better explains geographically structured phylogenies in legumes. *Phil. Trans. Roy. Soc. London B*, 359: 1509-1522.
- Lavin, M., Herendeen, P. S., & Wojciechowski, M. F. 2005. Evolutionary rates analysis of Leguminosae implicates a rapid diversification of lineages during the Tertiary. *Syst. Biol.* 54: 530-549.
- Lavin, M., Pennington, R. T., Hughes, C. E., Lewis, G. P., Delgado-Salinas, A., Dundo-de-Stefano, R., Queiroz, L. P. de, Cardoso, D., & Wojciechowski, M. F. 2018. DNA sequence variation among conspecific accessions of the legume *Coursetia caribaea* reveals geographically localized clades here ranked as species. *Systematic Botany* 43: 664-675.
- Legume Phylogeny Working Group (LPWG) [including D. Cardoso, L. P. de Queiroz, & M. F. Wojciechowski]. 2013. Toward a new classification system for legumes: progress report from the ILC6. *S. Afr. J. Bot.* 89: 3-9. doi:10.1016/j.sajb.2013.07.022.
- Legume Phylogeny Working Group (LPWG) [including D. Cardoso, L. P. de Queiroz, & M. F. Wojciechowski]. 2013. Legume phylogeny and classification in the 21st century: progress, prospects and lessons for other species-rich clades. *Taxon* 62: 217-248.
- Legume Phylogeny Working Group (LPWG) [including D. Cardoso, H. C. de Lima, L. P. de Queiroz, & M. F. Wojciechowski]. 2017. A new subfamily classification of the Leguminosae based on a taxonomically comprehensive phylogeny. *Taxon* 66: 44-77. doi.org/10.12705/661.
- Queiroz, L.P., Lewis, G.P., Wojciechowski, M.F., 2010. *Tabaroa*, a new genus of Leguminosae tribe Brongniartieae from Brazil. *Kew Bull.* 65: 189-203.
- Queiroz, L. P. de, Pastore, J. F. B., Cardoso, D., Snak, C., Lima, A. L. C., Gagnon, E., Vatanparast, M., Holland, A. E., & Egan, A. N. 2015. A multilocus phylogenetic analysis reveals the monophyly of a recircumscribed papilionoid legume tribe Diocleae with well-supported generic relationships. *Mol. Phylogenet. Evol.* 90: 1-19.
- Queiroz, L. P. de, W. São-Mateus, A. Delgado-Salinas, B. M. Torke, G. P. Lewis, O. Dorado, J. K. Ardley, M. F. Wojciechowski, and D. Cardoso. 2017. A molecular phylogeny reveals the Cuban enigmatic genus *Behaimia* as a new piece in the Brongniartieae puzzle of papilionoid legumes. *Mol. Phylogenet. & Evol.* 109: 191-202.
- Ramos, G., de Lima, H. C., Prenner, G., de Queiroz, L. P., Zartman, C. E., & Cardoso, D. 2016. Molecular systematics of the Amazonian genus *Aldina*, a phylogenetically enigmatic ectomycorrhizal lineage of papilionoid legumes. *Molec. Phyl. Evol.* 97: 11-18.
- Sabir J, Schwarz EN, Ellison N, Zhang J, Baeshen NA, Mutwakil M, Jansen RK, Ruhlman TA. 2014. Evolutionary and biotechnology implications of plastid genome variation in the inverted-repeat lacking clade of legumes. *Plt Biotechnol J* 12: 743–754.

- Sanderson, M. J. & Wojciechowski, M. F. 1996. Diversification rates in a temperate legume clade: Are there “so many species” of *Astragalus* (Fabaceae)? *Amer. J. Bot.* 83: 1488-1502.
- Saski C, Lee S, Daniell H, Wood T, Tomkins J, Kim H-G, Jansen RK. 2005. Complete chloroplast genome sequence of *Glycine max* and comparative analyses with other legume genomes. *Plant Mol Biol* 59: 309–322.
- Schwarz EN, Ruhlman T, Sabir JSM, Hajrah NH, Alharbi NS, Al-Malki AL, Bailey CD, Jansen RK. 2015. Plastid genomes reveal parallel inversions and multiple losses of *rps16* in papilionoids. *J Syst Evol* 53: 458–468.
- Schwarz EN, Ruhlman TA, Weng M-L, Khiyami MA, Sabir JMS, Hajarrah NH, Alharbi NS, Rabah SO, Jansen RK. 2017. Plastome-wide nucleotide substitution rates reveal accelerated rates in Papilionoideae and correlations with genome features across legume subfamilies. *J Mol Evol* 84: 187–203.
- Silva, M. J. da, Queiroz, L. P. de, Tozzi, A. M. G. A., Lewis, G. P., & Sousa, A. P. de. 2012. Phylogeny and biogeography of *Lonchocarpus* sensu lato and its allies in the tribe Millettieae (Leguminosae, Papilionoideae). *Taxon* 61: 93-108.
- Snak, C., Vatanparast, M., Silva, C., Lewis, G. P., Lavin, M., Kajita, T., de Queiroz, L. P. 2016. A dated phylogeny of the papilionoid legume genus *Canavalia* reveals recent diversification by a pantropical liana lineage. *Molec. Phyl. Evol* 98: 133-146.
- Steele KP, Wojciechowski MF. 2003. Phylogenetic analyses of tribes Trifolieae and Viciae, based on sequences of the plastid gene, *matK* (Papilionoideae: Leguminosae). In Klitgaard BB, Bruneau A.(eds). *Advances in legume systematics, part 10*, 355 – 370. Royal Botanic Gardens, Kew, UK.
- Steele KP, Ickert-Bond SM, Zarre S, Wojciechowski MF. 2010. Phylogeny and character evolution in *Medicago* (Leguminosae): Evidence from analyses of plastid *trnK/matK* and nuclear *GA3OX1* sequences. *Amer J Bot* 97: 1142–1155.
- Swanepoel, W., le Roux, M. M., Wojciechowski, M. F., & van Wyk, A. E., 2015. *Oberholzeria* (Fabaceae subfam. Faboideae), a new monotypic legume genus from Namibia. *PLoS ONE* 10: e0122080.
- Wojciechowski, M. F., Sanderson, M. J., Baldwin, B. G., & Donoghue, M. J. 1993. Monophyly of aneuploid *Astragalus*: Evidence from nuclear ribosomal DNA internal transcribed spacer sequences. *Amer. J. Bot.* 80: 711-722.
- Wojciechowski, M. F., Sanderson, M. J., & Hu, J.-M. 1999. Evidence on the monophyly of *Astragalus* (Fabaceae) and its major subgroups based on nuclear ribosomal DNA ITS and chloroplast DNA *trnL* intron data. *Syst. Bot.* 24: 409-437.
- Wojciechowski, M. F., Sanderson, M. J., & Hu, J.-M. 1999. Evidence on the monophyly of *Astragalus* (Fabaceae) and its major subgroups based on nuclear ribosomal DNA ITS and chloroplast DNA *trnL* intron data. *Syst. Bot.* 24: 409-437.
- Wojciechowski, M. F., M. J. Sanderson, K. P. Steele, and A. Liston. 2000. Molecular phylogeny of the “temperate herbaceous tribes” of papilionoid legumes: a supertree approach. In *Advances in Legume Systematics*, P. Herendeen and A. Bruneau (eds.), part 9, pp. 277-298. Royal Botanic Garden, Kew.
- Wojciechowski, M. F. 2003. Reconstructing the phylogeny of legumes (Leguminosae): an early 21st century perspective. In *Advances in Legume Systematics, Higher Level Systematics*, B. B. Klitgaard and A. Bruneau (eds.), part 10, pp. 5-35. Royal Botanic Gardens, Kew.

- Wojciechowski MF, Lavin M, Sanderson MJ. 2004. A phylogeny of legumes (Leguminosae) based on analysis of the plastid *matK* gene resolves many well-supported subclades within the family. *Amer J Bot* 91: 1846–1862.
- Wojciechowski, M. F. 2004. *Astragalus* (Fabaceae): A molecular phylogenetic perspective. *Brittonia* 57: 382-396.
- Wojciechowski, M. F., Sanderson Wojciechowski, M. F. 2013. Towards a new classification of Leguminosae: naming clades using non-Linnaean phylogenetic nomenclature. *S. Afr. J. Bot.* 89: 85-93.
- Zhang R., Wang Y.-H., Jin J.-J, Stull G.W., Bruneau A., Cardoso D., de Queiroz L.P., Moore M.J., Zhang S.D., Chen S.-Y., Wang J., Li D.-Z., Yi T.-S. 2020. Exploration of plastid phylogenomic conflict yields new insights into the deep relationships of Leguminosae. *Syst Biol*, <https://doi.org/10.1093/sysbio/syaa013>.